

100 Integrals

(Great for calc 1 and calc 2 students)

Video: <https://youtu.be/dgm4-3-lv3s>

@blackpenredpen

March 1st, 2019

Video transcription by jackw11111

$$\begin{aligned}
 \text{(Q1.) } & \int \tan^5 x \sec^3 x \, dx \\
 &= \int \tan^4 x \sec^2 x \tan x \sec x \, dx \\
 &= \int (u^2 - 1)^2 u^2 \, du \\
 &= \int (u^6 - 2u^4 + u^2) \, du \\
 &= \boxed{\frac{1}{7} \sec^7 x - \frac{2}{5} \sec^5 x + \frac{1}{3} \sec^3 x + C}
 \end{aligned}$$

Aside

$$\begin{aligned}
 \tan^4 x &= (\tan^2 x)^2 \\
 &= (\sec^2 x - 1)^2 \\
 \text{Let } u &= \sec x \\
 du &= \sec x \tan x \, dx
 \end{aligned}$$

$$\begin{aligned}
 \text{(Q2.) } & \int \frac{\cos 2x}{\sin x + \cos x} \, dx \\
 &= \int \frac{(\cos x - \sin x)(\cos x + \sin x)}{\sin x + \cos x} \, dx \\
 &= \int (\cos x - \sin x) \, dx \\
 &= \boxed{\sin x + \cos x + C}
 \end{aligned}$$

Aside

$$\begin{aligned}
 \cos(2x) &= \cos^2 x - \sin^2 x \\
 &= (\cos x - \sin x)(\cos x + \sin x)
 \end{aligned}$$

$$\begin{aligned}
 \text{(Q3.) } & \int \frac{x^2 + 1}{x^4 - x^2 + 1} \, dx \\
 &= \int \frac{1 + \frac{1}{x^2}}{x^2 - 1 + \frac{1}{x^2}} \, dx \\
 &= \int \frac{du}{u^2 + 1} \\
 &= \boxed{\tan^{-1} \left(x - \frac{1}{x} \right) + C}
 \end{aligned}$$

Aside

$$\begin{aligned}
 x^2 - 1 + \frac{1}{x^2} &= x^2 - 2 + \frac{1}{x^2} + 1 \\
 &= \left(x - \frac{1}{x} \right)^2 + 1 \\
 \text{Let } u &= x - \frac{1}{x} \\
 du &= \left(1 + \frac{1}{x^2} \right) dx
 \end{aligned}$$

$$\begin{aligned}
 \text{(Q4.) } & \int (x + e^x)^2 \, dx \\
 &= \int (x^2 + 2xe^x + e^{2x}) \, dx \\
 &= \boxed{\frac{1}{3}x^3 + 2xe^x - 2e^x + \frac{1}{2}e^{2x} + C}
 \end{aligned}$$

Aside

	<u>D</u>	<u>I</u>
+	2x	e ^x
-	2	e ^x
+	0	e ^x

$$\begin{aligned}
(Q5.) \int \csc^3 x \sec x \, dx &= \int \frac{1}{\sin^3 x \cos x} \, dx \\
&= \int \frac{\cancel{\sin^2 x} \cdot 1}{\cancel{\sin^2 x} \cos x} + \frac{\cancel{\cos^2 x}}{\sin^3 x \cancel{\cos x}} \, dx \\
&= \int \frac{\sin^2 x + \cos^2 x}{\sin x \cos x} + \frac{\cos x}{\sin^3 x} \, dx \\
&= \int \frac{\sin^2 x}{\sin x \cos x} + \frac{\cos^2 x}{\sin x \cos x} + \frac{\cos x}{\sin^3 x} \, dx \\
&= \int \frac{\sin x}{\cos x} \, dx + \int \frac{\cos x}{\sin x} \, dx + \int \frac{\cos x}{\sin^3 x} \, dx \\
&= \ln |\sec x| + \ln |\sin x| - \frac{1}{2 \sin^2 x} \\
&= \boxed{\ln |\tan x| - \frac{1}{2} \csc^2 x + C}
\end{aligned}$$

Aside

$$1 = \sin^2 x + \cos^2 x$$

$$\begin{aligned}
\ln |\sec x| + \ln |\sin x| &= \ln (|\sec x| \times |\sin x|) \\
&= \ln |\tan x|
\end{aligned}$$

$$\begin{aligned}
(Q6.) \int \frac{\cos x}{\sin^2 x - 5 \sin x - 6} \, dx &= \int \frac{1}{u^2 - 5u - 6} \, du \\
&= \int \frac{\frac{1}{7}}{u-6} + \frac{-\frac{1}{7}}{u+1} \, du \\
&= \frac{1}{7} \ln |u-6| - \frac{1}{7} \ln |u+1| \\
&= \boxed{\frac{1}{7} \ln |\sin x - 6| - \frac{1}{7} \ln |\sin x + 1| + C}
\end{aligned}$$

Aside

$$\text{Let } u = \sin x$$

$$du = \cos x \, dx$$

$$u^2 - 5u - 6 = (u-6)(u+1)$$

$$u - 6 = 0 \text{ when } u = 6$$

$$\text{sub } u = 6 \text{ into } u + 1 = 7$$

$$u + 1 = 0 \text{ when } u = -1$$

$$\text{sub } u = -1 \text{ into } u - 6 = -7$$

$$\begin{aligned}
(Q7.) \int \frac{1}{\sqrt{e^x}} \, dx &= \int \frac{1}{e^{\frac{x}{2}}} \, dx \\
&= \int e^{-\frac{x}{2}} \, dx \\
&= -2e^{-\frac{x}{2}} \\
&= \boxed{\frac{-2}{\sqrt{e^x}} + C}
\end{aligned}$$

Aside

$$\sqrt{n} = n^{\frac{1}{2}}$$

$$\int e^{ax} \, dx = \frac{1}{a} e^{ax} + C$$

$$\begin{aligned}
 \text{(Q8.) } & \int \frac{e^x \sqrt{e^x - 1}}{e^x + 3} dx \\
 &= \int \frac{\cancel{e^x} u}{u^2 + 1 + 3} \frac{2u}{\cancel{e^x}} du \\
 &= 2 \int \frac{u^2 + 4 - 4}{u^2 + 4} du \\
 &= 2 \left(\int 1 du - 4 \int \frac{1}{u^2 + 2^2} du \right) \\
 &= 2 \left(u - 4 \cdot \frac{1}{2} \tan^{-1} \left(\frac{u}{2} \right) \right) \\
 &= \boxed{2\sqrt{e^x - 1} - 4 \tan^{-1} \left(\frac{\sqrt{e^x - 1}}{2} \right) + C}
 \end{aligned}$$

Aside


$$\begin{aligned}
 \text{Let } u &= \sqrt{e^x - 1} \\
 du &= \frac{e^x}{2\sqrt{e^x - 1}} dx \\
 dx &= \frac{2\sqrt{e^x - 1}}{e^x} du \\
 u^2 + 1 &= e^x
 \end{aligned}$$

$$\begin{aligned}
 \text{(Q9.) } & \int \frac{1}{x + \sqrt{x}} dx \\
 &= \int \frac{1}{\sqrt{x}(\sqrt{x} + 1)} dx \\
 &= \int \frac{1}{\sqrt{x} \times u} \frac{2\sqrt{x}}{2\sqrt{x}} du \\
 &= \boxed{2 \ln(\sqrt{x} + 1) + C}
 \end{aligned}$$

Aside

$$\begin{aligned}
 \int \frac{1}{x + \sqrt{x}} dx &\neq \ln|x + \sqrt{x}| + C \\
 \text{Let } u &= \sqrt{x} + 1 \\
 du &= \frac{1}{2\sqrt{x}} dx \\
 dx &= 2\sqrt{x} du
 \end{aligned}$$

$$\begin{aligned}
 \text{(Q10.) } & \int_{-1}^5 |x - 3| dx \\
 &= 8 + 2 \\
 &= \boxed{10}
 \end{aligned}$$



$$\begin{aligned}
 \text{(Q11.) } & \int \frac{\sin x}{\sec^{2019} x} dx \\
 &= \int \cos^{2019} x \sin x dx \\
 &= \int u^{2019} \sin x - \frac{du}{\sin x} \\
 &= \boxed{-\frac{1}{2020} \cos^{2020} x + C}
 \end{aligned}$$

Aside

$$\begin{aligned}
 \text{Let } u &= \cos x \\
 du &= -\sin x dx \\
 dx &= -\frac{du}{\sin x}
 \end{aligned}$$

$$\begin{aligned}
 \text{(Q12.) } & \int \frac{x \sin^{-1} x}{\sqrt{1-x^2}} dx \\
 &= -\sin^{-1} x \sqrt{1-x^2} + \int 1 dx \\
 &= \boxed{-\sin^{-1} x \sqrt{1-x^2} + x + C}
 \end{aligned}$$

Aside

$$\begin{array}{r}
 D \\
 + \sin^{-1} x \\
 - \frac{1}{\sqrt{1-x^2}} \rightarrow \frac{x}{\sqrt{1-x^2}} \\
 \cdot
 \end{array}$$

$$\begin{aligned}
 \int u dv &= \int -\frac{1}{\sqrt{1-x^2}} x - \sqrt{1-x^2} dx \\
 &= \int 1 dx
 \end{aligned}$$

$$\begin{aligned}
 \text{(Q13.) } & \int \frac{2 \sin x}{\sin(2x)} dx \\
 &= \int \frac{2 \sin x}{2 \sin x \cos x} dx \\
 &= \int \sec x dx \\
 &= \int \frac{\sec x (\sec x + \tan x)}{\sec x + \tan x} dx \\
 &= \int \frac{1}{u} du \\
 &= \boxed{\ln |\sec x + \tan x| + C}
 \end{aligned}$$

Aside

$$\begin{aligned}
 \sin 2x &= 2 \sin x \cos x \\
 \text{Let } u &= \sec x + \tan x \\
 du &= \sec x \tan x + \sec^2 x dx \\
 &= \sec x (\tan x + \sec x) dx
 \end{aligned}$$

$$\begin{aligned}
 \text{(Q14.) } & \int \cos^2(2x) dx \\
 &= \frac{1}{2} \int (1 + \cos 4x) dx \\
 &= \frac{1}{2} \left(x + \frac{1}{4} \sin 4x \right) \\
 &= \boxed{\frac{1}{2}x + \frac{1}{8} \sin 4x + C}
 \end{aligned}$$

Aside

$$\begin{aligned}
 \cos^2 x &= \frac{1}{2}(1 + \cos(2x)) \\
 \cos^2 2x &= \frac{1}{2}(1 + \cos(4x))
 \end{aligned}$$

$$\begin{aligned}
 \text{(Q15.) } & \int \frac{1}{x^3 + 1} dx \\
 &= \frac{1}{3} \int \frac{1}{x+1} dx - \frac{1}{3} \int \frac{x-2}{x^2-x+1} dx \\
 &= \frac{1}{3} \int \frac{1}{x+1} dx - \frac{1}{3} \int \frac{2(x-1-3)}{x^2-x+1} dx \\
 &= \frac{1}{3} \int \frac{1}{x+1} dx - \frac{2}{3} \int \frac{x-4}{x^2-x+1} dx \\
 &= \frac{1}{3} \ln|x+1| - \frac{2}{3} \left(\frac{1}{2} \ln|x^2-x+1| + \frac{1}{\sqrt{3}} \tan^{-1} \left(\frac{x-\frac{1}{2}}{\frac{\sqrt{3}}{2}} \right) \right) \\
 &= \boxed{\frac{1}{3} \ln|x+1| - \frac{1}{6} \ln|x^2-x+1| + \frac{1}{\sqrt{3}} \tan^{-1} \left(\frac{2x}{\sqrt{3}} - \frac{1}{\sqrt{3}} \right) + C}
 \end{aligned}$$

Aside

$$\begin{aligned}
 (x^3 + 1) &= (x+1)(x^2-x+1) \\
 \frac{1}{(x+1)(x^2-x+1)} &= \frac{A}{x+1} + \frac{Bx+C}{x^2-x+1}
 \end{aligned}$$

$$x+1 = 0 \text{ when } x = -1$$

$$\text{substituting } x = -1 \text{ into } x^2 - x + 1 = 3$$

$$A = \frac{1}{3}$$

used $x = -1$, now we sub $x = 0$

$$1 = \frac{\frac{1}{3}}{0+1} + \frac{B \times 0 + C}{0^2 - 0 + 1}$$

$$1 = \frac{1}{3} + C$$

$$C = \frac{2}{3}$$

use $x = 1$

$$\frac{1}{(1+1)(1^2-1+1)} = \frac{\frac{1}{3}}{1+1} + \frac{B \times 1 + \frac{2}{3}}{1^2-1+1}$$

$$\frac{1}{2} = \frac{1}{6} + B + \frac{2}{3}$$

$$B = -\frac{1}{3}$$

$$\frac{d}{dx}(x^2 - x + 1) = 2x - 1$$

$$x^2 - x + 1 = x^2 - x + \frac{1}{4} + \frac{3}{4}$$

$$\begin{aligned}
 \text{(Q16.) } & \int x \sin^2 x \, dx \\
 &= \frac{1}{2} \int x (1 - \cos(2x)) \, dx \\
 &= \frac{1}{2} \left(\int x \, dx + \int -x \cos(2x) \, dx \right) \\
 &= \frac{1}{2} \left(\frac{1}{2} x^2 - \frac{1}{2} x \sin(2x) - \frac{1}{4} \cos(2x) \right) \\
 &= \boxed{\frac{1}{4} x^2 - \frac{1}{4} x \sin(2x) - \frac{1}{8} \cos(2x) + C}
 \end{aligned}$$

Aside

$$\begin{aligned}
 \sin^2 x &= \frac{1}{2} (1 - \cos(2x)) \\
 & \begin{array}{r} D \qquad I \\ + \quad -x \quad \cos^2 x \\ - \quad -1 \quad \frac{1}{2} \sin 2x \\ + \quad 0 \quad -\frac{1}{4} \cos 2x \end{array}
 \end{aligned}$$

$$\begin{aligned}
 \text{(Q17.) } & \int \left(x + \frac{1}{x} \right)^2 dx \\
 &= \int x^2 + 2 + x^{-2} \, dx \\
 &= \boxed{\frac{1}{3} x^3 + 2x - \frac{1}{x} + C}
 \end{aligned}$$

$$\begin{aligned}
 \text{(Q18.) } & \int \frac{3}{x^2 + 4x + 29} \, dx \\
 &= 3 \int \frac{1}{(x+2)^2 + 5^2} \, dx \\
 &= \boxed{\frac{3}{5} \tan^{-1} \left(\frac{x+2}{5} \right) + C}
 \end{aligned}$$

Aside

$$\begin{aligned}
 x^2 + 4x + 29 &= x^2 + 4x + 4 + 25 \\
 &= (x+2)^2 + 5^2
 \end{aligned}$$

$$\begin{aligned}
 \text{(Q19.) } & \int \cot^5 x \, dx \\
 &= \int \frac{\cos^5 x}{\sin^5 x} \, dx \\
 &= \int \frac{(1 - \sin^2 x)^2 \cos x}{\sin^5 x} \, dx \\
 &= \int \frac{(1 - u^2)^2}{u^5} \, du \\
 &= \int u^{-5} - 2u^{-3} + u^{-1} \, du \\
 &= \boxed{-\frac{1}{4} \csc^4 x + \csc^2 x + \ln |\sin x| + C}
 \end{aligned}$$

Aside

$$\begin{aligned}
 \cos^4 x &= (1 - \sin^2 x)^2 \\
 \text{Let } u &= \sin x \\
 du &= \cos x \\
 \frac{1}{\sin x} &= \csc x
 \end{aligned}$$

$$(Q20.) \int_{-1}^1 \frac{\tan x}{x^4 - x^2 + 1} dx$$

$$= \boxed{0}$$

Aside

$$\frac{\text{odd } f(x)}{\text{even } f(x)} = \text{odd } f(x)$$

$$\int_{-a}^a \text{odd } f(x) dx = 0$$

$$(Q21.) \int \sin^3 x \cos^2 x dx$$

$$= \int (1 - \cos^2 x) \cos^2 x \sin x dx$$

$$= - \int (1 - u^2) u^2 du$$

$$= - \int (u^2 - u^4) du$$

$$= \boxed{-\frac{1}{3} \cos^3 x + \frac{1}{5} \cos^5 x + C}$$

Aside

$$\sin^3 x = \sin^2 x \times \sin x$$

$$= (1 - \cos^2 x) \times \sin x$$

Let $u = \cos x$
 $du = -\sin x dx$

$$(Q22.) \int \frac{1}{x^2 \sqrt{x^2 + 1}} dx$$

$$= \int \frac{x^{-3}}{\sqrt{1 + x^{-2}}} dx$$

$$= -\frac{1}{2} \int \frac{1}{\sqrt{u}} du$$

$$= -\frac{1}{2} \int u^{-\frac{1}{2}} du$$

$$= \boxed{-\sqrt{1 + \frac{1}{x^2}} + C}$$

Aside

$$x^2 \sqrt{x^2 + 1} = x^2 \sqrt{x^2 \left(\frac{x^2}{x^2} + \frac{1}{x^2} \right)}$$

$$= x^2 x \sqrt{1 + \frac{1}{x^2}}$$

$$= \frac{x^{-3}}{\sqrt{1 + x^{-2}}}$$

Let $u = 1 + x^{-2}$
 $du = 2x^{-3} dx$

$$(Q23.) \int \sin x \sec x \tan x dx$$

$$= \int \tan^2 x dx$$

$$= \int (\sec^2 x - 1) dx$$

$$= \boxed{\tan x - x + C}$$

Aside

$$\sin x \sec x = \frac{\sin x}{\cos x}$$

$$= \tan x$$

$$\begin{aligned}
 (Q24.) \int \sec^3 x \, dx & \\
 &= \sec x \tan x - \int \sec x \tan^2 x \, dx \\
 &= \sec x \tan x - \int \sec x (\sec^2 x - 1) \, dx \\
 &= \sec x \tan x - \int \sec^3 x \, dx + \int \sec x \, dx
 \end{aligned}$$

$$2 \int \sec^3 x \, dx = \sec x \tan x + \ln |\sec x + \tan x|$$

$$= \boxed{\frac{1}{2} \sec x \tan x + \frac{1}{2} \ln |\sec x + \tan x| + C}$$

Aside

$$\begin{array}{r}
 + \quad \overset{D}{\sec x} \quad \overset{I}{\sec^2 x} \\
 - \quad \sec x \tan x \quad \tan x \\
 \cdot
 \end{array}$$

$$\tan^2 x = \sec^2 x - 1$$

$$\begin{aligned}
 (Q25.) \int \frac{1}{x\sqrt{9x^2-1}} \, dx & \\
 &= \int \frac{1}{\frac{1}{3} \sec \theta \sqrt{\sec^2 \theta - 1}} \, dx \\
 &= \int \frac{1}{\frac{1}{3} \sec \theta \tan \theta} \frac{1}{3} \sec \theta \tan \theta \, d\theta \\
 &= \int 1 \, d\theta \\
 &= \theta \\
 &= \boxed{\sec^{-1}(3x) + C}
 \end{aligned}$$

Aside

$$9x^2 = (3x)^2$$

$$\text{Let } 3x = \sec \theta$$

$$x = \frac{1}{3} \sec \theta$$

$$dx = \frac{1}{3} \sec \theta \tan \theta \, d\theta$$

$$\sec^2 \theta = \tan^2 \theta + 1$$

$$\sec^2 \theta - 1 = \tan^2 \theta$$

$$\sqrt{\sec^2 \theta - 1} = \sqrt{\tan^2 \theta}$$

$$\sqrt{\sec^2 \theta - 1} = \tan \theta$$

$$\sec \theta = 3x$$

$$\theta = \sec^{-1} 3x$$

$$\begin{aligned}
 (Q26.) \int \cos \sqrt{x} \, dx & \\
 &= 2 \int u \cos u \, du \\
 &= 2u \sin u + 2 \cos u \\
 &= \boxed{2 \sqrt{x} \sin \sqrt{x} + 2 \cos \sqrt{x} + C}
 \end{aligned}$$

Aside

$$\text{Let } u = \sqrt{x}$$

$$x = u^2$$

$$dx = 2u \, du$$

$$\begin{array}{r}
 \overset{D}{2u} \quad \overset{I}{\cos u} \\
 + \quad 2u \quad \cos u \\
 - \quad 2 \quad \sin u \\
 + \quad 0 \quad -\cos u
 \end{array}$$

$$\begin{aligned}
 (Q27.) \int \csc x \, dx &= \int \frac{\csc x (\csc x - \cot x)}{(\csc x - \cot x)} \, dx \\
 &= \int \frac{1}{u} \, du \\
 &= \boxed{\ln |\csc x - \cot x| + C}
 \end{aligned}$$

Aside

$$\text{Let } u = \csc x - \cot x$$

$$du = -\csc x \cot x + \cos^2 x \, dx$$

$$\begin{aligned}
 (Q28.) \int \sqrt{x^2 + 4x + 13} \, dx &= \int \sqrt{(x+2)^2 + 3^2} \, dx \\
 &= \int \sqrt{(3 \tan \theta)^2 + 3^2} (3 \sec^2 \theta) \, d\theta \\
 &= \int 3 \sqrt{\tan^2 \theta + 1} (3 \sec^2 \theta) \, d\theta \\
 &= 9 \int \sec^3 \theta \, d\theta \\
 &= 9 \left(\frac{1}{2} \sec \theta \tan \theta + \frac{1}{2} \ln |\sec \theta + \tan \theta| \right) \\
 &= \frac{9}{2} \frac{\sqrt{x^2 + 4x + 13}}{3} \times \frac{(x+2)}{3} + \frac{9}{2} \ln \left| \frac{\sqrt{x^2 + 4x + 13}}{3} + \frac{x+2}{3} \right| + C_1 \\
 &= \boxed{\frac{1}{2}(x+2)\sqrt{x^2 + 4x + 13} + \frac{9}{2} \ln(\sqrt{x^2 + 4x + 13} + (x+2)) + C_2}
 \end{aligned}$$

Aside

$$\sqrt{x^2 + 4x + 13} = \sqrt{x^2 + 4x + 4 + 9}$$

$$= \sqrt{(x+2)^2 + 3^2}$$

$$\text{Let } x+2 = 3 \tan \theta$$

$$dx = 3 \sec^2 \theta \, d\theta$$


$$\sqrt{\tan^2 \theta + 1} = \sqrt{\sec^2 \theta}$$

$$= \sec \theta$$

$$\text{recall } \int \sec^3 x \, dx$$

$$= \frac{1}{2} \sec x \tan x + \frac{1}{2} \ln |\sec x + \tan x|$$

$$\tan \theta = \frac{x+2}{3}$$



$$C_2 = 3 + C_1$$

$$\begin{aligned}
 (Q29.) \int e^{2x} \cos x \, dx &= e^{2x} \sin x + 2e^{2x} \cos x - 4 \int e^{2x} \cos x \, dx + 4 \int e^{2x} \cos x \, dx \\
 \int e^{2x} \cos x \, dx &= \boxed{\frac{1}{5} e^{2x} \sin x + \frac{2}{5} e^{2x} \cos x + C}
 \end{aligned}$$

Aside

$$\begin{array}{r}
 D \quad I \\
 + \quad e^{2x} \cos x \\
 - \quad 2e^{2x} \sin x \\
 + \quad 4e^{2x} \cos x
 \end{array}$$

$$(Q30.) \int_3^5 (x-3)^9 dx$$

$$= \int_{u=0}^{u=2} u^9 du$$

$$= \left[\frac{1}{10} u^{10} \right]_0^2$$

$$= \frac{1}{10} 2^{10} - \frac{1}{10} 0^{10}$$

$$= \frac{1}{10} 2^{10} - \frac{1}{10} 0^{10}$$

$$= \boxed{\frac{512}{5}}$$

Aside

$$\text{Let } u = x - 3$$

$$du = dx$$

$$(Q31.) \int \frac{1}{\sqrt{x-x^{3/2}}} dx$$

$$= \int \frac{1}{\sqrt{x}\sqrt{1-\sqrt{x}}} dx$$

$$= \int \frac{-2\sqrt{x}}{\sqrt{x}\sqrt{u}} du$$

$$= -2 \int 2u^{-1/2} + 1 = \frac{1}{2} du$$

$$= -4u^{1/2}$$

$$= \boxed{-4\sqrt{1-\sqrt{x}} + C}$$

Aside

$$\sqrt{x-x^{3/2}} = \sqrt{x(1-x^{1/2})}$$

$$= \sqrt{x}\sqrt{1-\sqrt{x}}$$

$$\text{Let } u = 1 - \sqrt{x}$$

$$du = \frac{-1}{2\sqrt{x}} dx$$

$$dx = -2\sqrt{x} du$$

$$(Q32.) \int \frac{1}{\sqrt{x-x^2}} dx$$

$$= \int \frac{1}{\sqrt{x}\sqrt{1-x}} dx$$

$$= \int \frac{1}{\sqrt{x}\sqrt{1-(\sqrt{x})^2}} dx$$

$$= \int \frac{1}{\sqrt{x}\sqrt{1-u^2}} 2\sqrt{x} du$$

$$= 2 \sin^{-1}(u)$$

$$= \boxed{2 \sin^{-1}(\sqrt{x}) + C}$$

Aside

$$\text{Let } u = \sqrt{x}$$

$$du = \frac{1}{2\sqrt{x}} dx$$

$$dx = 2\sqrt{x} du$$

$$\begin{aligned}
 \text{(Q33.) } & \int e^{2 \ln x} dx \\
 &= \int e^{\ln(x^2)} dx \\
 &= \int x^2 dx \\
 &= \boxed{\frac{1}{3}x^3 + C}
 \end{aligned}$$

Aside

$$y \ln(x) = \ln(x^y)$$

$$\begin{aligned}
 \text{(Q34.) } & \int \frac{\ln x}{\sqrt{x}} dx \\
 &= 2\sqrt{x} \ln x - 2 \int \frac{1}{\sqrt{x}} dx \\
 &= \boxed{2\sqrt{x} \ln(x) - 4\sqrt{x} dx + C}
 \end{aligned}$$

Aside

$$\begin{array}{r}
 D \quad I \\
 + \ln x \quad x^{-\frac{1}{2}} \\
 - \quad -\frac{1}{x} \quad 2\sqrt{x}
 \end{array}$$

$$\begin{aligned}
 -\frac{1}{x} \times 2\sqrt{x} &= -2 \frac{\sqrt{x}}{x} \\
 &= -2 \frac{\cancel{\sqrt{x}}}{\sqrt{x} \sqrt{x}} \\
 &= -\frac{2}{\sqrt{x}}
 \end{aligned}$$

$$\begin{aligned}
 \text{(Q35.) } & \int \frac{1}{e^x + e^{-x}} dx \\
 &= \int \frac{1(e^x)}{e^x(e^x + e^{-x})} dx \\
 &= \int \frac{e^x}{(e^x)^2 + 1} dx \\
 &= \int \frac{u}{u^2 + 1} du \\
 &= \boxed{\tan^{-1}(e^x) + C}
 \end{aligned}$$

Aside

$$\begin{aligned}
 \text{Let } u &= e^x \\
 du &= e^x dx
 \end{aligned}$$

$$\begin{aligned}
 \text{(Q36.) } & \int \log_2 x \, dx \\
 &= \int \frac{\ln x}{\ln 2} \, dx \\
 &= \frac{1}{\ln 2} \int \ln x \, dx \\
 &= \frac{1}{\ln 2} (x \ln x - x) \, dx \\
 &= \frac{x \ln x}{\ln 2} - \frac{x}{\ln 2} \, dx \\
 &= \boxed{x \log_2(x) - \frac{x}{\ln 2} + C}
 \end{aligned}$$

Aside

$$\begin{array}{r}
 D \quad I \\
 + \quad \ln x \quad 1 \\
 - \int \frac{1}{x} \cdot x \, dx
 \end{array}$$

$$\begin{aligned}
 \text{(Q37.) } & \int x^3 \sin(2x) \, dx \\
 &= \boxed{-\frac{1}{2}x^3 \cos(2x) + \frac{3}{4}x^2 \sin(2x) + \frac{3}{4}x \cos(2x) - \frac{3}{8} \sin(2x) + C}
 \end{aligned}$$

Aside

$$\begin{array}{r}
 D \quad I \\
 + \quad x^3 \quad \sin(2x) \\
 - \quad 3x^2 \quad -\frac{1}{2} \cos 2x \\
 + \quad 6x \quad \frac{1}{4} \sin 2x \\
 - \quad 6 \quad \frac{1}{8} \cos 2x \\
 + \quad 0 \quad \frac{1}{16} \sin 2x
 \end{array}$$

$$\begin{aligned}
 \text{(Q38.) } & \int x^2 \sqrt[3]{1+x^3} \, dx \\
 &= \frac{1}{3} \int u^{\frac{1}{3}} \, du \\
 &= \frac{1}{\cancel{3}} \frac{3}{4} u^{\frac{4}{3}} \, du \\
 &= \frac{1}{4} \sqrt[3]{(1+x^3)^4} + C \\
 &= \boxed{\frac{1}{4}(1+x^3) \sqrt[3]{1+x^3} + C}
 \end{aligned}$$

Aside

$$\begin{aligned}
 \text{Let } u &= 1 + x^3 \\
 du &= 3x^2 \, dx \\
 \frac{1}{3} du &= x^2 \, dx
 \end{aligned}$$

$$\begin{aligned}
(Q39.) \int \frac{1}{(x^2 + 4)^2} dx &= \int \frac{1}{(4 \sec^2 \theta)^2} 2 \sec^2 \theta d\theta \\
&= \int \frac{1}{(4 \sec^2 \theta)(4 \sec^2 \theta)} 2 \sec^2 \theta d\theta \\
&= \frac{1}{8} \int \frac{1}{\sec^2 \theta} d\theta \\
&= \frac{1}{8} \int \cos^2 \theta d\theta \\
&= \frac{1}{8} \frac{1}{2} \int (1 + \cos(2\theta)) d\theta \\
&= \frac{1}{16} (\theta + \frac{1}{2} \sin(2\theta)) \\
&= \frac{1}{16} \theta + \frac{1}{16} \sin \theta \cos \theta \\
&= \frac{1}{16} \tan^{-1} \left(\frac{x}{2} \right) + \frac{1}{16} \frac{x}{\sqrt{x^2 + 4}} \frac{2}{\sqrt{x^2 + 4}} \\
&= \boxed{\frac{1}{16} \tan^{-1} \left(\frac{x}{2} \right) + \frac{1}{8} \frac{x}{x^2 + 4} + C}
\end{aligned}$$

Aside

$$\text{Let } x = 2 \tan \theta$$

$$x^2 + 4 = 4 \tan^2 \theta + 4$$

$$x^2 + 4 = 4(\tan^2 \theta + 1)$$

$$x^2 + 4 = 4 \sec^2 \theta$$


$$dx = 2 \sec^2 \theta d\theta$$

$$\cos^2 \theta = \frac{1}{2}(1 + \cos(2\theta))$$

$$\sin 2\theta = 2 \sin \theta \cos \theta$$


$$\frac{1}{2} \sin 2\theta = \sin \theta \cos \theta$$

$$\text{remember } \tan \theta = \frac{x}{2}$$



$$\begin{aligned}
(Q40.) \int_1^2 \sqrt{x^2 - 1} dx &= \boxed{\sqrt{3} - \frac{\cosh^{-1}(2)}{2}} \\
\text{or} & \\
&= \boxed{\sqrt{3} - \frac{\sinh^{-1}(\sqrt{3})}{2}}
\end{aligned}$$

Aside



$$\text{When } x = 2, y = \sqrt{2^2 - 1} = \sqrt{3}$$

$$\text{Area of triangle} = \frac{1}{2} \times 2 \times \sqrt{3} - \frac{t}{2}$$

$$\cosh t = 2, \sinh t = \sqrt{3}$$

$$t = \cosh^{-1}(2), t = \sinh^{-1}(\sqrt{3})$$

$$\text{Area} = \sqrt{3} - \frac{\cosh^{-1}(2)}{2} \text{ or } \sqrt{3} - \frac{\sinh^{-1}(\sqrt{3})}{2}$$

$$\begin{aligned}
 \text{(Q41.) } \int \sinh x \, dx &= \int \frac{e^x - e^{-x}}{2} \, dx \\
 &= \frac{1}{2}(e^x + e^{-x}) \\
 &= \boxed{\cosh x + C}
 \end{aligned}$$

$$\begin{aligned}
 \text{(Q42.) } \int \sinh^2 x \, dx &= \frac{1}{2} \int -1 + \cosh(2x) \, dx \\
 &= \frac{1}{2}(-x + \frac{1}{2} \sinh(2x)) \\
 &= \boxed{-\frac{1}{2}x + \frac{1}{4} \sinh(2x) + C}
 \end{aligned}$$

$$\begin{aligned}
 \text{(Q43.) } \int \sinh^3 x \, dx &= \int \sinh^2 \sinh x \, dx \\
 &= \int (\cosh^2 x - 1) \sinh x \, dx \\
 &= \int (u^2 - 1) \, du \\
 &= \frac{1}{3}u^3 - u \\
 &= \boxed{\frac{1}{3} \cosh^3 x - \cosh x + C}
 \end{aligned}$$

Aside

$$\begin{aligned}
 \sinh x &= \frac{e^x - e^{-x}}{2} \\
 \sinh^2 x &= \left(\frac{e^x - e^{-x}}{2} \right)^2 \\
 &= \frac{1}{4}(e^x - e^{-x})^2 \\
 &= \frac{1}{4}(e^{2x} - 2 + e^{-2x}) \\
 &= \frac{1}{4} \times (-2) + \frac{1}{4}(e^{2x} + e^{-2x}) \\
 &= -\frac{1}{2} + \frac{1}{2 \times 2}(e^{2x} + e^{-2x}) \\
 &= -\frac{1}{2} + \frac{1}{2} \left(\frac{e^{2x} + e^{-2x}}{2} \right) \\
 &= -\frac{1}{2} + \frac{1}{2} \cosh(2x)
 \end{aligned}$$

Aside

$$\begin{aligned}
 \cosh^2 x - \sinh^2 x &= 1 \\
 \sinh^2 x &= \cosh^2 x - 1 \\
 \text{Let } u &= \cosh x \\
 du &= \sinh x \, dx
 \end{aligned}$$

$$(Q44.) \int \frac{1}{\sqrt{x^2 + 1}} dx$$

$$= \boxed{\sinh^{-1} x + C}$$

Aside

$$\frac{d}{dx}(\sin^{-1} x) = \frac{1}{\sqrt{1 - x^2}}$$

$$\frac{d}{dx}(\sinh^{-1} x) = \frac{1}{\sqrt{1 + x^2}}$$

$$(Q45.) \int \ln(x + \sqrt{1 + x^2}) dx$$

$$= x \sinh^{-1} x - \int \frac{x}{\sqrt{1 + x^2}} dx$$

$$= \boxed{x \sinh^{-1} x - \sqrt{1 + x^2} + C}$$

Aside

$$\begin{array}{r} D \qquad I \\ + \sinh^{-1}(x) - \frac{1}{x} \\ - \int \frac{1}{\sqrt{1+x^2}} dx \end{array}$$

$$(Q46.) \int \tanh x dx$$

$$= \int \frac{\sinh x}{\cosh x} dx$$

$$= \int \frac{du}{u}$$

$$= \boxed{\ln |\cosh x| + C}$$

Aside

Let $u = \cosh(x)$
 $du = \sinh(x) dx$

$$(Q47.) \int \operatorname{sech} x dx$$

$$= \int \frac{1 \times \cosh x}{\cosh x \cosh x} dx$$

$$= \int \frac{\cosh x}{1 + \sinh^2 x} dx$$

$$= \int \frac{1}{1 + u^2} du$$

$$= \boxed{\tan^{-1}(\sinh x) + C}$$

Aside

Let $u = \sinh x$
 $du = \cosh x dx$

$$(Q48.) \int \tanh^{-1} x \, dx$$

$$= x \tanh^{-1} x - \int \frac{x}{1-x^2} \, dx$$

$$= x \tanh^{-1} x + \frac{1}{2} \ln|1-x^2| + C$$

Aside

$$+ \overset{D}{\sinh^{-1}(x)} - \overset{I}{\int \frac{1}{1+x^2} \, dx}$$

$$(Q49.) \int \sqrt{\tanh x} \, dx$$

$$= \int u \times \frac{2u}{1-u^4} \, du$$

$$= \int \frac{u^2 + 1 + u^2 - 1}{(1+u^2)(1-u^2)} \, du$$

$$= \int \frac{\cancel{u^2+1} + \cancel{-(1-u^2)}}{(1+u^2)(1-u^2)} \, du$$

$$= \tanh^{-1}(\sqrt{\tanh x}) - \tan^{-1}(\sqrt{\tanh x}) + C$$

Aside

$$\text{Let } u = \sqrt{\tanh x}$$

$$x = \tanh^{-1}(u^2)$$

$$dx = \frac{2u}{1-(u^2)^2} \, du$$


$$1-u^4 = (1+u^2)(1-u^2)$$

Aside

$$(Q50.) \int_0^5 [x] \, dx$$

$$= 1 + 2 + 3 + 4$$

$$= \boxed{10}$$



$$\begin{aligned}
 \text{(Q51.) } & \int \sec^6 x \, dx \\
 &= \int (\tan^2 x + 1)^2 \sec^2 x \, dx \\
 &= \int (u^2 + 1)^2 \, du \\
 &= \int (u^4 + 2u^2 + 1) \, du \\
 &= \boxed{\frac{1}{5} \tan^5 x + \frac{2}{3} \tan^3 x + \tan x + C}
 \end{aligned}$$

Aside
 $u = \tan x$
 $du = \sec^2 x \, dx$

$$\begin{aligned}
 \text{(Q52.) } & \int \frac{1}{(5x-2)^4} \, dx \\
 &= \frac{1}{5} \int \frac{1}{u^4} \, du \\
 &= \frac{1}{5} \int -\frac{1}{3} u^{-4+1} \, du \\
 &= \\
 &= \boxed{-\frac{1}{15} \frac{1}{(5x+2)^3} + C}
 \end{aligned}$$

Aside
Let $u = 5x - 2$
 $du = 5 \, dx$
 $\frac{1}{5} du = dx$

$$\begin{aligned}
 \text{(Q53.) } & \int \ln(1+x^2) \, dx \\
 &= x \ln(1+x^2) - 2 \int \frac{1+x^2-1}{1+x^2} \, dx \\
 &= \boxed{x \ln(1+x^2) - 2x + 2 \tan^{-1}(x) + C}
 \end{aligned}$$

Aside

$$\begin{aligned}
 \text{(Q54.) } & \int \frac{1}{x^4+x} \, dx \\
 &= \int \frac{x^{-4}}{1+x^{-3}} \, dx \\
 &= \int \frac{x^{-4}}{u} \frac{du}{-3x^{-4}} \\
 &= \boxed{-\frac{1}{3} \ln|1+x^{-3}| + C}
 \end{aligned}$$

Aside
 $x^4 + x = x^4(1+x^{-3})$
Let $u = 1+x^{-3}$
 $du = -3x^{-4} \, dx$
 $dx = \frac{du}{-3x^{-4}}$

$$\begin{aligned}
 (Q55.) \int \frac{1 - \tan x}{1 + \tan x} dx &= \int \frac{1 - \frac{\sin x}{\cos x}}{1 + \frac{\sin x}{\cos x}} dx \\
 &= \int \frac{\frac{\cos x - \sin x}{\cos x}}{\frac{\cos x + \sin x}{\cos x}} dx \\
 &= \int \frac{\cos x - \sin x}{\cos x + \sin x} \times \frac{\cos x}{\cos x} dx \\
 &= \int \frac{\cos x - \sin x}{\cos x + \sin x} dx \\
 &= \int \frac{du}{u} \\
 &= \boxed{\ln |\cos x + \sin x| + C}
 \end{aligned}$$

Aside

$$\text{Let } u = \cos x + \sin x$$

$$du = \cos x - \sin x \, dx$$

$$\begin{aligned}
 (Q56.) \int x \sec x \tan x \, dx &= \boxed{x \sec x - \ln |\sec x + \tan x| + C}
 \end{aligned}$$

Aside

$$\begin{array}{r}
 D \qquad \qquad I \\
 + \ x \qquad \searrow \sec x \tan x \\
 - \ 1 \qquad \searrow \sec x \\
 \hline
 + \ 0 \qquad \ln |\sec x + \tan x|
 \end{array}$$

$$\begin{aligned}
 (Q57.) \int \sec^{-1} x \, dx &= x \sec^{-1} x - \int \frac{1}{x\sqrt{x^2-1}} dx \\
 &= x \sec^{-1} x - \int \frac{1}{\sqrt{\sec^2 \theta - 1}} \sec \theta \tan \theta \, d\theta \\
 &= x \sec^{-1} x - \int \frac{1}{\tan \theta} \sec \theta \tan \theta \, d\theta \\
 &= x \sec^{-1} x - \int \sec \theta \, d\theta \\
 &= x \sec^{-1} x - \ln |\sec \theta + \tan \theta| \\
 &= \boxed{x \sec^{-1} x - \ln |x + \sqrt{x^2 - 1}| + C}
 \end{aligned}$$


Aside

$$\begin{array}{r}
 D \qquad \qquad I \\
 + \ \sec^{-1} x \qquad 1 \\
 - \ \frac{1}{x\sqrt{x^2-1}} \qquad x \\
 \hline
 \end{array}$$

$$\text{Let } x = \sec \theta$$

$$dx = \sec \theta \tan \theta \, d\theta$$

$$\sec^2 \theta - 1 = \tan^2 \theta$$



$$\tan \theta = \sqrt{x^2 - 1}$$

$$\begin{aligned}
 \text{(Q58.) } & \int \frac{1 - \cos x}{1 + \cos x} dx \\
 &= \int \frac{2 \sin^2\left(\frac{x}{2}\right)}{2 \cos^2\left(\frac{x}{2}\right)} dx \\
 &= \int \tan^2\left(\frac{x}{2}\right) dx \\
 &= \int (\sec^2\left(\frac{x}{2}\right) - 1) dx \\
 &= \boxed{2 \tan\left(\frac{x}{2}\right) - x + C}
 \end{aligned}$$

Aside

$$\cos^2 \frac{x}{2} = \frac{1}{2}(1 + \cos x)$$

$$\sin^2 \frac{x}{2} = \frac{1}{2}(1 - \cos x)$$

$$\begin{aligned}
 \text{(Q59.) } & \int x^2 \sqrt{x+4} dx \\
 &= \int (u-4)^2 \sqrt{u} du \\
 &= \int (u^2 - 8u + 16) u^{\frac{1}{2}} du \\
 &= \int \left(\frac{2}{7} u^{\frac{5}{2}+1} - \frac{2}{5} 8 u^{\frac{3}{2}+1} + 16 \times \frac{2}{3} u^{\frac{1}{2}+1} \right) du \\
 &= \boxed{\frac{2}{7}(x+4)^{\frac{7}{2}} - \frac{16}{5}(x+4)^{\frac{5}{2}} + \frac{2}{3}(x+4)^{\frac{3}{2}} + C}
 \end{aligned}$$

Aside


$$\text{Let } u = x + 4$$

$$du = dx$$

$$x = u - 4$$

$$\begin{aligned}
 \text{(Q60.) } & \int_{-1}^1 \sqrt{4-x^2} dx \\
 &= 2 \int_0^1 \sqrt{4-x^2} dx \\
 &= 2 \left(\frac{1}{2} \times 1 \times \sqrt{3} + \frac{1}{2} 2^2 \times \frac{\pi}{6} \right) \\
 &= \boxed{\sqrt{3} + \frac{2\pi}{3}}
 \end{aligned}$$

Aside



$$\theta = \sin^{-1}\left(\frac{1}{2}\right) = \frac{\pi}{6}$$

$$\text{Area of Sector} = \frac{1}{2} r^2 \theta$$

$$\begin{aligned}
(Q61.) \int \sqrt{x^2 + 4x} \, dx &= \int \sqrt{(x+2)^2 - 2^2} \, dx \\
&= \int \sqrt{4 \sec^2 \theta - 4} \, 2 \sec \theta \tan \theta \, d\theta \\
&= 4 \int \tan^2 \theta \sec \theta \, d\theta \\
&= 4 \int (\sec^2 \theta - 1) \sec \theta \, d\theta \\
&= 4 \int \sec^3 \theta - \sec \theta \, d\theta \\
&= 4 \left(\frac{1}{2} \sec \theta \tan \theta + \frac{1}{2} \ln |\sec \theta + \tan \theta| - \ln |\sec \theta + \tan \theta| \right) \\
&= 4 \left(\frac{1}{2} \sec \theta \tan \theta - \frac{1}{2} \ln |\sec \theta + \tan \theta| \right) \\
&= 2 \frac{x+2}{2} \frac{\sqrt{x^2+4x}}{2} - 2 \ln \left| \frac{x+2}{2} + \frac{\sqrt{x^2+4x}}{2} \right| + C_1 \\
&= \boxed{\frac{1}{2}(x+2)\sqrt{x^2+4x} - 2 \ln |(x+2) + \sqrt{x^2+4x}| + C_2}
\end{aligned}$$

Aside


$$x^2 + 4x + 4 - 4 = (x+2)^2 - 2^2$$

$$\text{Let } x+2 = 2 \sec \theta$$

$$dx = 2 \sec \theta \tan \theta \, d\theta$$

$$4(\sec^2 \theta - 1) = 4 \tan^2 \theta$$

$$\sec \theta = \frac{x+2}{2}$$



$$\begin{aligned}
(Q62.) \int x^2 e^{x^3} \, dx &= \frac{1}{3} \int 3x^2 e^{x^3} \, dx \\
&= \boxed{\frac{1}{3} e^{x^3} + C}
\end{aligned}$$

$$\begin{aligned}
(Q63.) \int x^3 e^{x^2} \, dx &= \frac{1}{2} x^2 e^{x^2} - \frac{1}{2} \int 2x e^{x^2} \, dx \\
&= \boxed{\frac{1}{2} x^2 e^{x^2} - \frac{1}{2} e^{x^2} + C}
\end{aligned}$$

Aside

$$\begin{array}{r}
D \quad I \\
+ \quad x^2 \quad x e^{x^2} \\
- \quad 2x \quad \frac{1}{2} e^{x^2}
\end{array}$$

$$(Q64.) \int \tan x \ln(\cos x) dx$$

$$= \int \cancel{\tan x} u \frac{du}{\cancel{-\tan x}}$$

$$= - \int u du$$

$$= \boxed{-\frac{1}{2} (\ln(\cos x))^2 + C}$$

Aside

$$\text{Let } u = \ln(\cos x)$$

$$du = \frac{-\sin x}{\cos x} dx$$

$$du = -\tan x dx$$

$$dx = \frac{du}{-\tan x}$$

$$(Q65.) \int \frac{1}{x^3 - 4x^2} dx$$

$$= \int \frac{-\frac{1}{16}}{x} + \frac{-\frac{1}{4}}{x^2} + \frac{\frac{1}{16}}{x-4} dx$$

$$= \boxed{-\frac{1}{16} \ln|x| + \frac{1}{4x} + \frac{1}{16} \ln|x-4| + C}$$

Aside

$$x^3 - 4x^2 = x^2(x-4)$$

$$\frac{1}{x^3 - 4x^2} = \frac{Ax+B}{x^2} + \frac{C}{x-4}$$

$$\frac{1}{x^3 - 4x^2} = \frac{Ax}{x^2} + \frac{B}{x^2} + \frac{C}{x-4}$$

$$= \frac{A}{x} + \frac{B}{x^2} + \frac{C}{x-4}$$

When $x=4$

$$C = \frac{1}{16}$$

When $x=0$

$$B = -\frac{1}{4}$$

When $x=1$

$$-\frac{1}{3} = A - \frac{1}{4} - \frac{1}{48}$$

$$-\frac{16}{48} = A - \frac{12}{48} - \frac{1}{48}$$

$$A = -\frac{1}{16}$$

$$(Q66.) \int \sin x \cos(2x) dx$$

$$= - \int (2u^2 - 1) du$$

$$= - \left(\frac{2u^3}{3} - u \right)$$

$$= \boxed{-\frac{2}{3} \cos^3 x + \cos x + C}$$

Aside

$$\cos(2x) = 2\cos^2 x - 1$$

$$\text{Let } u = \cos x$$

$$du = -\sin x dx$$

$$\begin{aligned}
(Q67.) \int 2^{\ln x} dx &= \int (e^{\ln 2})^{\ln x} dx \\
&= \int (e^{\ln x})^{\ln 2} dx \\
&= \int (x)^{\ln 2} dx \\
&= \int \frac{1}{(\ln 2 + 1)} x^{(\ln 2) + 1} dx \\
&= \frac{1}{1 + \ln 2} x^{1 + \ln x} \\
&= \frac{1}{1 + \ln 2} x \times x^{\ln x} \\
&= \boxed{\frac{x 2^{\ln x}}{1 + \ln 2} + C}
\end{aligned}$$

$$\begin{aligned}
(Q68.) \int \sqrt{1 + \cos(2x)} dx &= \int \sqrt{2 \cos^2 x} dx \\
&= \sqrt{2} \int \cos x dx \\
&= \boxed{\sqrt{2} \sin x + C}
\end{aligned}$$

Aside

$$2 \cos^2 x = 1 + \cos(2x)$$

$$\begin{aligned}
(Q69.) \int \frac{1}{1 + \tan x} dx &= \frac{1}{2} \int \frac{1 - \tan x + 1 + \tan x}{1 + \tan x} dx \\
&= \frac{1}{2} \int \frac{1 - \tan x}{1 + \tan x} + \frac{1 + \tan x}{1 + \tan x} dx \\
&= \boxed{\frac{1}{2} \ln |\cos x + \sin x| + \frac{1}{2} x + C}
\end{aligned}$$

Aside

$$\text{recall } \int \frac{1 - \tan x}{1 + \tan x} = \ln |\cos x + \sin x| \text{ from Q55}$$

$$\begin{aligned}
 \text{(Q70.) } & \int_{\frac{1}{e}}^e \frac{\sqrt{1 - (\ln x)^2}}{x} dx \\
 &= \int_{-1}^1 \frac{\sqrt{1 - u^2}}{\cancel{x}} \cancel{x} du \\
 &= \boxed{\frac{\pi}{2}}
 \end{aligned}$$


Aside

Let $u = \ln x$

$$du = \frac{1}{x} dx$$

When $x = \frac{1}{e}, u = -1$

When $x = e, u = 1$



$$\begin{aligned}
 \text{Area} &= \frac{\pi r^2}{2} \\
 &= \frac{\pi}{2}
 \end{aligned}$$

$$\begin{aligned}
 \text{(Q71.) } & \int \frac{1}{\sqrt[3]{x} + 1} dx \\
 &= \int \frac{1}{u} 3(u-1)^2 du \\
 &= 3 \int \frac{(u-1)^2}{u} du \\
 &= 3 \int \frac{u^2 - 2u + 1}{u} du \\
 &= 3 \int \left(u - 2 + \frac{1}{u}\right) du \\
 &= 3\left(\frac{1}{2}u^2 - 2u + \ln|u|\right) \\
 &= \boxed{\frac{3}{2}(\sqrt[3]{x} + 1)^2 - 6(\sqrt[3]{x} + 1) + \ln|\sqrt[3]{x} + 1| + C}
 \end{aligned}$$

Aside

Let $u = \sqrt[3]{x} + 1$

$x = (u - 1)^3$

$dx = 3(u - 1)^2$

$$\begin{aligned}
 (Q72.) \int \frac{1}{\sqrt[3]{x+1}} dx & \\
 &= \frac{3}{2} \int u^{-\frac{1}{3}+1} du \\
 &= \frac{3}{2} (x+1)^{\frac{2}{3}} \\
 &= \boxed{\frac{3}{2} \sqrt[3]{(x+1)^2} + C}
 \end{aligned}$$

Aside

$$\text{Let } u = x + 1$$

$$du = dx$$

$$\begin{aligned}
 (Q73.) \int (\sin x + \cos x)^2 dx & \\
 &= \int \sin^2 x + 2 \sin x \cos x + \cos^2 x dx \\
 &= \int (1 + \sin(2x)) dx \\
 &= \boxed{x - \frac{1}{2} \cos(2x) + C}
 \end{aligned}$$

Aside

$$\sin^2 x + \cos^2 x = 1$$

$$2 \sin x \cos x = \sin 2x$$

$$\begin{aligned}
 (Q74.) \int 2x \ln(1+x) dx & \\
 &= x^2 \ln(1+x) - \int \frac{x^2}{1+x^2} dx \\
 &= x^2 \ln(1+x) - \int \left(x - 1 + \frac{1}{x+1} \right) dx \\
 &= \boxed{x^2 \ln(1+x) - \frac{1}{2}x^2 + x - \ln(x+1) + C}
 \end{aligned}$$

Aside

$$\begin{array}{r}
 D \quad I \\
 + \ln(1+x) \quad 2x \\
 - \frac{1}{1+x} \quad \frac{x^2}{x^2}
 \end{array}$$

$$\begin{aligned}
 &= x + 1 \sqrt{x^2} \\
 &\quad - (x^2 + x) \\
 &\quad \quad -x \\
 &\quad \quad \quad -(-x-1) \\
 &\quad \quad \quad \quad 1
 \end{aligned}$$

$$\begin{aligned}
(Q75.) \int \frac{1}{x(1 + \sin^2(\ln x))} dx \\
&= \int \frac{\frac{1}{\cos^2 u}}{\frac{1}{\cos^2 u} + \frac{\sin^2(u)}{\cos^2 u}} du \\
&= \int \frac{\sec^2 u}{\sec^2 u + \tan^2 u} du \\
&= \int \frac{\sec^2 u}{2 \tan^2 x + 1} du \\
&= \frac{1}{2 w^2 + 1} dw \\
&= \frac{1}{(\sqrt{2}w)^2 + 1} dw \\
&= \frac{1}{\sqrt{2}} \tan^{-1}(\sqrt{2}w) \\
&= \boxed{\frac{1}{\sqrt{2}} \tan^{-1}(\sqrt{2} \tan(\ln x)) + C}
\end{aligned}$$

Aside

$$\text{Let } u = \ln x$$

$$du = \frac{1}{x} dx$$

$$\sec^2 x = \tan^2 x + 1$$

$$\text{Let } w = \tan u$$

$$dw = \sec^2 u du$$

$$\begin{aligned}
(Q76.) \int \sqrt{\frac{1-x}{1+x}} dx \\
&= \int \sqrt{\frac{1-x(1-x)}{1+x(1-x)}} dx \\
&= \int \frac{1-x}{\sqrt{1-x^2}} dx \\
&= \int \frac{1}{\sqrt{1-x^2}} dx - \int \frac{x}{\sqrt{1-x^2}} dx \\
&= \boxed{\sin^{-1} x + \sqrt{1-x^2} + C}
\end{aligned}$$

$$\begin{aligned}
(Q77.) \int x^{\frac{x}{\ln x}} dx \\
&= \int (e^{\ln x})^{\frac{x}{\ln x}} dx \\
&= \int e^x dx \\
&= e^x \\
&= \boxed{x^{\frac{x}{\ln x}} + C}
\end{aligned}$$

Aside

$$\text{Let } u = \sqrt{x}$$

$$x = u^2$$

$$dx = 2u \, du$$

$$du = -\frac{1}{2\sqrt{x}}$$

$$\begin{array}{r}
 D \quad I \\
 + \sin^{-1} u \quad 2u \\
 - \frac{1}{\sqrt{1-u^2}} \rightarrow u^2
 \end{array}$$

$$(Q78.) \int \sin^{-1}(\sqrt{x}) \, dx$$

$$= \int 2u \sin^{-1} u \, du$$

$$= u^2 \sin^{-1} u - \int \frac{u^2}{\sqrt{1-u^2}} \, du$$

$$\boxed{x \sin^{-1} \sqrt{x} - \frac{1}{2} \sin^{-1} \sqrt{x} + \frac{1}{2} \sqrt{x} \sqrt{1-x} + C}$$

$$- \int \frac{u^2}{\sqrt{1-u^2}} \, du$$

$$\text{Let } u = \sin \theta$$

$$du = \cos \theta \, d\theta$$

$$- \int \frac{\sin^2 \theta}{\cos \theta} \cos \theta \, d\theta$$


$$= -\frac{1}{2} \int 1 - \cos 2\theta \, d\theta$$

$$= -\frac{1}{2} \left(\theta - \frac{1}{2} \sin 2\theta \right)$$

$$= -\frac{1}{2} \theta + \frac{1}{2} \sin \theta \cos \theta$$

$$\sin \theta = \frac{u}{1}$$

$$\theta = \sin^{-1} u$$



$$(Q79.) \int \tan^{-1} x \, dx$$

$$= x \tan^{-1} x - \int \frac{x}{1+x^2} \, dx$$

$$= x \tan^{-1} x - \frac{1}{2} \ln(1+x^2)$$

$$\boxed{x \tan^{-1} x - \frac{1}{2} \ln(1+x^2) + C}$$

Aside

$$\begin{array}{r}
 D \quad I \\
 + \tan^{-1} x \quad 1 \\
 - \frac{1}{1+x^2} \rightarrow x
 \end{array}$$

$$\begin{aligned}
 \text{(Q80.) } \int_0^5 f(x) dx, \text{ where } f(x) &= \begin{cases} 10 & \text{if } x \leq 2 \\ 3x^2 - 2 & \text{if } x > 2 \end{cases} dx \\
 &= \int_0^2 10 dx + \int_2^5 (3x^2 - 2) dx \\
 &= 20 + [x^3 - 2x]_2^5 \\
 &= 20 + [5^3 - 10 - 2^3 + 4] \\
 &= \boxed{131}
 \end{aligned}$$

$$\begin{aligned}
 \text{(Q81.) } \int \frac{\sin(\frac{1}{x})}{x^3} dx \\
 &= \boxed{\frac{\cos \frac{1}{x}}{x} - \sin(\frac{1}{x}) + C}
 \end{aligned}$$

Aside

$$\begin{array}{r}
 D \quad I \\
 + \quad \frac{1}{x^3} \quad \frac{\sin \frac{1}{x}}{x^2} \\
 - \quad \frac{-1}{x^2} \quad \cos \frac{1}{x} \\
 \hline
 \end{array}$$

$$\begin{aligned}
 \text{(Q82.) } \int \frac{x-1}{x^4-1} dx \\
 &= \int \frac{\frac{1}{2}}{x+1} dx - \frac{1}{2} \int \frac{x}{x^2+1} dx + \frac{1}{2} \int \frac{1}{x^2+1} dx \\
 &= \boxed{\frac{1}{2} \ln|x+1| - \frac{1}{4} \ln(x^2+1) + \frac{1}{2} \tan^{-1} x + C}
 \end{aligned}$$

Aside

$$\begin{aligned}
 x^4 - 1 &= (x-1)(x+1)(x^2+1) \\
 \frac{1}{(x-1)(x+1)(x^2+1)} &= \frac{A}{x+1} + \frac{Bx+C}{x^2+1} \\
 A &= \frac{1}{2} \\
 \text{using } x=0, C &= \frac{1}{2} \\
 \text{using } x=1, B &= -\frac{1}{2}
 \end{aligned}$$

$$\begin{aligned}
\text{(Q83.) } & \int \sqrt{1 + \left(x - \frac{1}{4x}\right)^2} dx \\
&= \int \sqrt{1 + x^2 - \frac{1}{2} + \frac{1}{16x^2}} dx \\
&= \int \sqrt{x^2 + \frac{1}{2} + \frac{1}{16x^2}} dx \\
&= \int \sqrt{\left(x + \frac{1}{4x}\right)^2} dx \\
&= \int x + \frac{1}{4x} dx \\
&= \boxed{\frac{1}{2}x^2 + \frac{1}{4} \ln|x| + C}
\end{aligned}$$

$$\begin{aligned}
\text{(Q84.) } & \int \frac{e^{\tan x}}{1 - \sin^2 x} dx \\
&= \int \sec^2 x e^{\tan x} dx \qquad \text{Aside} \\
& \qquad \qquad \qquad 1 - \sin^2 x = \cos^2 x \\
&= \boxed{e^{\tan x} + C}
\end{aligned}$$

$$\begin{aligned}
\text{(Q85.) } & \int \frac{\tan^{-1} x}{x^2} dx \\
&= -\frac{\tan^{-1} x}{x} + \int \frac{1}{x(1+x^2)} dx \\
&= -\frac{\tan^{-1} x}{x} + \int \frac{1}{x \times x^2(x^2+1)} dx \\
&= -\frac{\tan^{-1} x}{x} - \frac{1}{2} \int \frac{-2x^{-3}}{x^2+1} dx \\
&= \boxed{-\frac{\tan^{-1} x}{x} - \frac{1}{2} \ln(x^2+1) + C}
\end{aligned}$$

Aside

$$\begin{array}{r}
D \qquad \qquad I \\
+ \tan^{-1} x \quad \frac{1}{x^2} \\
- \frac{1}{1+x^2} \quad - \frac{1}{x} \\
\hline
\end{array}$$

$$\begin{aligned}
\text{(Q86.) } & \int \frac{\tan^{-1} x}{1+x^2} dx \\
&= \int u du \qquad \text{Aside} \\
&= \frac{1}{2} u^2 \qquad \text{Let } u = \tan^{-1} x \\
&= \boxed{\frac{1}{2} (\tan^{-1} x)^2 + C} \qquad du = \frac{1}{1+x^2} dx \\
& \qquad \qquad \qquad x = \tan u
\end{aligned}$$

$$\begin{aligned}
 \text{(Q87.) } \int (\ln x)^2 dx & \\
 &= x(\ln x)^2 - \int 2 \ln x dx \\
 &= x(\ln x)^2 - (2x \ln x - \int 2 dx) \\
 &= \boxed{x(\ln x)^2 - 2x \ln x + 2x + C}
 \end{aligned}$$

Aside

$$\begin{array}{r}
 D \quad I \\
 + \quad (\ln x)^2 \quad 1 \\
 - \quad 2 \ln x \quad x \\
 \hline
 \end{array}$$

$$\begin{array}{r}
 D \quad I \\
 + \quad 2 \ln x \quad 1 \\
 - \quad \frac{2}{x} \quad x \\
 \hline
 \end{array}$$

$$\begin{aligned}
 \text{(Q88.) } \int \frac{\sqrt{x^2+4}}{x^2} dx & \\
 &= \int \frac{2 \sec \theta}{4 \tan^2 \theta} 2 \sec^2 \theta d\theta \\
 &= \int \frac{1}{\cos^2 \theta} \times \frac{\cos^2 \theta}{\sin^2 \theta} d\theta \\
 &= \int \left(\frac{\sin^2 \theta}{\cos \theta \sin^2 \theta} + \frac{\cos^2 \theta}{\cos \theta \sin^2 \theta} \right) d\theta \\
 &= \ln |\sec \theta + \tan \theta| - \csc \theta \\
 &= \ln \left(\frac{\sqrt{x^2+4}}{2} + \frac{x}{2} \right) - \frac{\sqrt{x^2+4}}{x} + C_1 \\
 &= \boxed{\ln(\sqrt{x^2+4} + x) - \frac{\sqrt{x^2+4}}{x} + C_2}
 \end{aligned}$$

Aside

$$\text{Let } x = 2 \tan \theta$$

$$dx = 2 \sec^2 \theta d\theta$$


$$\sqrt{(2 \tan \theta)^2 + 4} = \sqrt{4(\tan^2 \theta + 1)}$$

$$= \sqrt{4(\sec^2 \theta)}$$

$$= 2 \sec \theta$$

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$\tan \theta = \frac{x}{2}$$



$$\begin{aligned}
 \text{(Q89.) } \int \frac{\sqrt{x+4}}{x} dx & \\
 &= \int \frac{u}{u^2-4} 2u du \\
 &= \int 2 + \frac{8}{u^2-4} du \\
 &= \int 2 + \frac{8}{(u-2)(u+2)} du \\
 &= \int 2 + \frac{2}{u-2} + \frac{-2}{u+2} du \\
 &= 2u + 2 \ln |u-2| - 2 \ln |u+2| \\
 &= \boxed{2\sqrt{x-4} + 2 \ln \left| \frac{\sqrt{x-4}-2}{\sqrt{x-4}+2} \right| + C}
 \end{aligned}$$

Aside

$$\text{Let } u = \sqrt{x+4}$$

$$u^2 = x+4$$

$$u^2 - 4 = x$$

$$dx = 2u du$$

$$= \frac{2}{u^2-4} 2u^2$$

$$= \frac{-(2u^2-8)}{u^2-4} = 8$$

$$(Q90.) \int_0^{\frac{\pi}{2}} \frac{\sin^3 x}{\sin^3 x + \cos^3 x} dx$$

$$= \boxed{\frac{\pi}{4}}$$

$$(Q91.) \int \frac{x}{1+x^4} dx$$

$$= \frac{1}{2} \int \frac{2x}{1+(x^2)^2} dx$$

$$= \boxed{\frac{1}{2} \tan^{-1}(x^2) + C}$$

$$(Q92.) \int e^{\sqrt{x}} dx$$

$$= \int 2ue^u du$$

$$= \boxed{2\sqrt{x}e^{\sqrt{x}} - 2e^{\sqrt{x}} + C}$$

Aside

$$\text{Let } u = \sqrt{x}$$

$$u^2 = x$$

$$2u du = dx$$

D	I	
+	2u	↘
-	2	↘
+	0	↘

$$(Q93.) \int \frac{1}{\csc^3 x} dx$$

$$= \int \sin^3 x dx$$

$$= \int \sin^2 x \sin x dx$$

$$= \int (1 - \cos^2 x) \sin x dx$$

$$= - \int (1 - u^2) du$$

$$= -\left(u - \frac{1}{3}u^3\right)$$

$$= \boxed{\frac{1}{3} \cos^3 x - \cos x + C}$$

Aside

$$\text{Let } u = \cos x$$

$$du = -\sin x dx$$

$$\begin{aligned}
 \text{(Q94.) } & \int \frac{\sin^{-1} x}{\sqrt{1-x^2}} dx \\
 &= \int \frac{u \cos u}{\cos u} du \\
 &= \frac{1}{2} u^2 \\
 &= \boxed{\frac{1}{2} (\sin^{-1} x)^2 + C}
 \end{aligned}$$

Aside

$$\text{Let } u = \sin^{-1} x$$

$$x = \sin u$$

$$dx = \cos u du$$

$$\sqrt{1 - \sin^2 u} = \cos u$$

$$\begin{aligned}
 \text{(Q95.) } & \int \sqrt{1 + \sin(2x)} dx \\
 &= \int \sqrt{\sin^2 x + 2 \sin x \cos x + \cos^2 x} dx \\
 &= \int \sqrt{(\sin x + \cos x)^2} dx \\
 &= \int \sin x + \cos x dx \\
 &= \boxed{-\cos x + \sin x + C}
 \end{aligned}$$

$$\begin{aligned}
 \text{(Q96.) } & \int \sqrt[4]{x} dx \\
 &= \int \frac{4}{5} x^{\frac{1}{4} + 1} dx \\
 &= \frac{4}{5} x^{\frac{5}{4}} \\
 &= \frac{4}{5} \sqrt[4]{x^5} \\
 &= \boxed{\frac{4}{5} x \sqrt[4]{x} + C}
 \end{aligned}$$

$$\begin{aligned}
 \text{(Q97.) } & \int \frac{1}{1 + e^x} dx \\
 &= \int \frac{1 + e^x - e^x}{1 + e^x} dx \\
 &= \int 1 dx - \int \frac{e^x}{1 + e^x} dx \\
 &= \boxed{x - \ln(1 + e^x) + C}
 \end{aligned}$$

$$\begin{aligned}
(Q98.) \int \sqrt{1+e^x} dx &= \int u \frac{2u}{u^2-1} du \\
&= \int 2 + \frac{2}{u^2-1} du \\
&= \int 2 + \frac{2}{(u-1)(u+1)} du \\
&= \int 2 + \frac{1}{u-1} - \frac{1}{u+1} du \\
&= 2u + \ln \left| \frac{u-1}{u+1} \right| \\
&= \boxed{2\sqrt{1+e^x} + \ln\left(\frac{\sqrt{1+e^x}-1}{\sqrt{1+e^x}+1}\right) + C}
\end{aligned}$$

Aside

$$\begin{aligned}
\text{Let } u &= \sqrt{1+e^x} \\
du &= \frac{e^x}{2\sqrt{1+e^x}} dx \\
dx &= \frac{2\sqrt{1+e^x}}{e^x} du \\
u^2 - 1 &= e^x \\
&= \frac{2}{u^2 - 1} \frac{2u^2}{2} \\
&= \frac{-(2u^2 - 2)}{2}
\end{aligned}$$

$$\begin{aligned}
(Q99.) \int \frac{\sqrt{\tan x}}{\sin(2x)} dx &= \int \frac{u}{2 \sin x \cos x} dx \\
&= \int \frac{u}{2 \sin x \cos x} \times \frac{2u}{\sec^2 x} du \\
&= \int \frac{u^2}{\sin x \cos x} \times \frac{\cos^2 x}{1} du \\
&= \int u^2 \cot x du \\
&= \int u^2 \frac{1}{u^2} du \\
&= \int 1 du \\
&= u \\
&= \boxed{\sqrt{\tan x} + C}
\end{aligned}$$

Aside

$$\begin{aligned}
\text{Let } u &= \sqrt{\tan x} \\
du &= \frac{\sec^2 x}{2\sqrt{\tan x}} dx \\
dx &= \frac{2\sqrt{\tan x}}{\sec^2 x} du \\
\tan x &= u^2 \\
\frac{1}{\tan x} &= \frac{1}{u^2}
\end{aligned}$$

$$\begin{aligned}
(Q100.) \int_0^{\frac{\pi}{2}} \frac{1}{1 + \sin x} dx &= \int \frac{1(1 - \sin x)}{1 + \sin x(1 - \sin x)} dx \\
&= \int \frac{1 - \sin x}{\cos^2 x} dx \\
&= \int (\sec^2 x - \sec x \tan x) dx \\
&= [\tan x - \sec x]_0^{\frac{\pi}{2}} \\
&= \left[\frac{\sin x}{\cos x} - \frac{1}{\cos x} \right]_0^{\frac{\pi}{2}} \\
&= \left[\frac{\sin x - 1(\sin x + 1)}{\cos x(\sin x + 1)} \right]_0^{\frac{\pi}{2}} \\
&= \left[\frac{-\cancel{\cos x}}{\cancel{\cos x}(1 + \sin x)} \right]_0^{\frac{\pi}{2}} \\
&= \left[\frac{-\cos x}{1 + \sin x} \right]_0^{\frac{\pi}{2}} \\
&= \boxed{1}
\end{aligned}$$

$$\begin{aligned}
(Q101.) \int \left(\frac{\sin x}{x} + \ln x \cos x \right) dx &= \boxed{\sin x \ln x + C}
\end{aligned}$$